

**Soil Erosion Enhancement Activity – SOE03 - Continuous no-till (organic system)**



**Enhancement Description**

This enhancement is for using a continuous no-till, strip till or direct seeding method of planting throughout the planned rotation on an organic farm. High residue levels are maintained by including high residue-producing crops, or by low residue crops followed by a cover crop in the rotation. Termination of all cover crops is accomplished using non-chemical methods, such as flail mowing, roller crimper and frost kill. No herbicides are used for weed control.

**Land Use Applicability**

Cropland (that is certified as organic or on conventional acreage that is being transitioned for organic certification).

**Benefits**

Use of continuous no-till, strip till or direct seeding leaves high levels of crop residue that can reduce erosion by wind and water up to 90%, increase soil organic matter, and control weeds. Mechanically terminating cover crops using a flail mower or roller crimper can eliminate the use of herbicides, thereby reducing potential offsite water quality problems while leaving the soil undisturbed.

**Criteria**

Implementation of this enhancement requires the use of continuous no-till, strip till or direct seeding on all crops during the planned rotation that is part of an organic system plan. The no-till, strip till or direct seeding system must incorporate the following activities:

1. Rotations that include only high residue producing crops
  - a. No cover crop required
  - b. Use only crops that produce high residue levels throughout the rotation, e.g. corn, wheat
  - c. Maintain a minimum of 90% residue cover on the soil surface after no-till, strip till or direct seed planting all crops
2. Rotations that include low residue crops
  - a. Use a cover crop after ALL low residue crops, e.g. vegetables, cotton, soybeans
  - b. Plant cover crops using a no till system



United States Department of Agriculture  
Natural Resources Conservation Service

2010 Ranking Period 2

- c. Maintain a minimum of 90% residue cover on the soil after no-till planting all crops
  - d. Use warm-season cover crops between spring and late summer crops or prior to late summer vegetable production
3. Additional Criteria
- a. All residues must be uniformly distributed over the entire field
  - b. No full-width tillage is permitted regardless of the depth of the tillage operation
  - c. Field(s) must have a soil loss at or below the tolerance (T) level for wind and/or water erosion for the crop rotation and a Soil Tillage Intensity Rating (STIR) of 30 or less for each planted crop or cover crop in the rotation
4. No Herbicides are used for weed control
5. Termination of all cover crops is accomplished using non-chemical methods, such as flail mowing, roller crimper and frost kill.

#### **Documentation**

1. Planned crop rotation showing cover crops that will be used after low residue crops
2. Planting method used for each crop in the rotation (no-till, strip till, direct seeding)
3. List of all other potential ground disturbing farming operations
4. Method of cover crop termination, e.g. flail mowing, roller crimper
5. Dates for farming operations
6. Map showing fields, acreage
7. Photographs of planted crops

## Indiana CSP Enhancement Supplemental Information

### **SOE03 – Continuous No-till (Organic System)**

- The **STIR** value is the **Soil Tillage Intensity Rating**. The RUSLE 2 software program utilizes the speed, depth, surface disturbance percent and tillage type parameters to calculate a tillage intensity rating for the system used in growing a crop or a rotation. STIR ratings tend to show the differences in the degree of soil disturbance between systems. The greater the disturbance, the higher STIR.

### **Crop Residue and Cover Crop List**

<b>Low Residue Crops <u>1/</u></b>	<b>High Residue Crops <u>2/</u></b>	<b>Cover Crops <u>3/</u></b>
Soybeans, Corn silage, Beans (dry edible), Cucumbers, Green peas, Green beans, Lima beans, Melons, Peppers, Pumpkins, Squash, Strawberries, Tomatoes, Vegetables, or similar crops	Corn (grain, sweet), Wheat, Barley Milo, Oats, Millet, Popcorn, Rye, Sorghum, Sorghum-Sudangrass Hybrids, Triticale	Radish- (forage, oilseed), Alsike Clover, Annual Ryegrass, Barley, Buckwheat, Canola/rape, Cowpeas, Crabgrass (red river), Crimson Clover, Field Peas/winter peas, Hairy Vetch, Mustards, Oats, Red Clover, Rye, Sorghum-Sudangrass Hybrids, Sweet clover, Triticale, Turnips, Wheat, Woollypod Vetch
<u>1/</u> Full/partial-season crops managed with cover crops	<u>2/</u> Full-season crops managed to leave 50 percent or more residue cover. Not harvested for silage or biomass.	<u>3/</u> Cover to be established early enough in growing season to provide adequate cover.